

On the gauge coupling unification

E.K. Loginov*

Department of Physics, Ivanovo State University, Ermaka St. 39, Ivanovo, 153025, Russia

We consider a quark-lepton symmetry model of unification of the strong and electromagnetic interactions. The model has the gauge group $SU(4) \times U(1)_Y$ and the minimal Higgs structure consisting of one complex quartet of scalar particles. The spontaneous breakdown of the gauge group to $SU(3)_c \times U(1)_{em}$ due to nonzero vacuum expectation value of the Higgs quartet provides the simplest realization of the Higgs mechanism which generates masses for gauge bosons, and masses to quarks and leptons. Using the embedding of the gauge group to $SU(5)$, we study the evolution of coupling constants and find a connection of the couplings with the gauge couplings of the standard model.

Despite the apparent striking success of the standard model (SM), there are a number of properties that this model does not explain, including the quantization of the electric charges of elementary particles, the ratios of the values of the respective standard-model gauge couplings g_s , g , and g' , and the interconnected manner in which quark and lepton contributions to gauge anomalies cancel each other (separately for each generation). For the SM to be consistent the Higgs boson mass should be relatively light $M_H \leq 1$ TeV. To explain the smallness of the Higgs boson mass some delicate cancellation is required that is rather nontrivial “fine tuning” or gauge hierarchy problem. Other nontrivial problem is that the SM can't predict the fermion masses, which vary over at least five orders of magnitude (fermion problem). These and other deficiencies of the SM motivated the effort to construct theories with higher unification of gauge symmetries.

One of the possible variants of such theory can be induced by the possible four color quark-lepton symmetry of Pati-Salam type treating leptons as the fourth color [1] (see also [2]). In the present paper we propose a quark-lepton symmetry model of unification of the strong and electromagnetic interactions. The model to be discussed here is based on the $SU(4) \times U(1)_Y$ group as the minimal group containing the four-color symmetry of quarks and leptons. Noting that $SU(4) \times U(1)_Y$ as a flavor group has been considered before in the literature (see e.g. [3]). However, in such papers as a rule possible extensions of the Glashow-Weinberg-Salam theory is studied. Whereas we construct an analogue of the electroweak theory whose symmetry group $SU(4) \times U(1)_Y$ breaks down to $SU(3)_c \times U(1)_{em}$.

We start by the following observation [4]. The gauge group $SU(5)$ has two inequivalent maximal breaking patterns:

$$SU(5) \rightarrow SU(3)_c \times SU(2) \times U(1)_Y, \quad (1)$$

$$SU(5) \rightarrow SU(4) \times U(1)_Y. \quad (2)$$

In the first case, we obtain the Georgi-Glashow $SU(5)$ -model [5]. We shall study the second case of the breaking patterns. As in the Georgi-Glashow model, we place the first generation of fermions in the $\mathbf{5}^* + \mathbf{10}$ representation

of $SU(5)$ in the following way:

$$\begin{pmatrix} d^{c1} \\ d^{c2} \\ d^{c3} \\ e \\ -\nu \end{pmatrix}_L, \quad \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & u^{c3} & -u^{c2} & u_1 & d_1 \\ -u^{c3} & 0 & u^{c1} & u_2 & d_2 \\ u^{c2} & -u^{c1} & 0 & u_3 & d_3 \\ -u_1 & -u_2 & -u_3 & 0 & e^c \\ -d_1 & -d_2 & -d_3 & -e^c & 0 \end{pmatrix}_L \quad (3)$$

Now we note that the simplest representations of $SU(5)$ have the following $SU(4) \times U(1)$ content:

$$\mathbf{5}^* = 1(-2) + 4^*(1/2), \quad (4)$$

$$\mathbf{10} = 4(3/2) + 6(-1), \quad (5)$$

$$\mathbf{24} = 1(0) + 15(0) + 4(-2) + 4^*(2). \quad (6)$$

These representations uniquely (to within the chirality of fermions) determine multiplets of fermions and gauge fields. Using (4) and (5), we get the following multiplets of spinor fields:

$$e_L, \quad \psi_L = \begin{pmatrix} d^{c1} \\ d^{c2} \\ d^{c3} \\ -\nu \end{pmatrix}_L, \quad \psi_R = \begin{pmatrix} u^{c1} \\ u^{c2} \\ u^{c3} \\ -e \end{pmatrix}_R, \quad (7)$$

and

$$\chi = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & u^{c3} & -u^{c2} & d_1 \\ -u^{c3} & 0 & u^{c1} & d_2 \\ u^{c2} & -u^{c1} & 0 & d_3 \\ -d_1 & -d_2 & -d_3 & 0 \end{pmatrix}_L. \quad (8)$$

Using the adjoint representation (6) we find multiplets of the vector fields: sixteen massless gauge bosons of $SU(4) \times U(1)$, six gauge bosons $X'_\mu{}^\alpha$ and $X'_{\mu\alpha}$ with the charges $\pm 4/3$, and two gauge bosons $Y'_\mu{}^{\pm}$. The latter eight gauge bosons obtain the masses via the Higgs mechanism.

Now we write out the Lagrangian of $SU(4) \times U(1)_Y$ -model. As in the Glashow-Weinberg-Salam model, it can be divided to five parts:

$$\mathcal{L} = \mathcal{L}_{YM} + \mathcal{L}_{HYM} + \mathcal{L}_{SH} + \mathcal{L}_f + \mathcal{L}_{Yuk}, \quad (9)$$

where

$$\mathcal{L}_{YM} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} - \frac{1}{4}G_{\mu\nu} G^{\mu\nu} \quad (10)$$

is the Yang-Mills Lagrangian without matter fields,

$$\mathcal{L}_{HYM} = (D_\mu \Phi)^+ (D^\mu \Phi) \quad (11)$$

is the Lagrangian describing the Higgs quartet interaction with $SU(4) \times U(1)_Y$ gauge fields,

$$\mathcal{L}_{SH} = \mu^2 \Phi^+ \Phi - \lambda (\Phi^+ \Phi)^2 \quad (12)$$

is the Lagrangian describing Higgs doublet self-interaction,

$$\mathcal{L}_f = \bar{\psi}^i i \gamma^\mu (D_\mu \psi)^i + \bar{\chi}_{ij} i \gamma^\mu (D_\mu \chi)_{ij} + \bar{e}_L i \gamma^\mu D_\mu e_L \quad (13)$$

is the Lagrangian describing the interaction of fermions with gauge fields, and

$$\mathcal{L}_{Yuk} = -f_e \bar{\psi}_R^i \Phi^i e_L - f_u \bar{\psi}_R^i \Phi^{ijk} \chi_{jk} - f_d \bar{\psi}_L^i \Phi^{ijk} \chi_{jk} + h.c. \quad (14)$$

is the Lagrangian describing the Yukawa interaction of the fermions with Higgs quartet.

Here the covariant derivatives is given by

$$D_\mu \Psi = \left(\partial_\mu - ig \frac{A_a}{2} A_\mu^a - ig' \frac{Y}{2} B_\mu \right) \Psi, \quad (15)$$

where values of A_a dependent on the choice of the multiplet Ψ . In particular, if $\Psi = e_L$, then all $A_a = 0$. If $\Psi = \chi$, then $A_p = 0$ as $p = 1, \dots, 7$ and

$$A_s = \frac{1}{\sqrt{6}} \begin{pmatrix} \lambda_s & 0 \\ 0 & 0 \end{pmatrix} \quad (16)$$

as $s = 8, \dots, 15$. Here λ_s are the usual Gell-Mann matrices for $SU(3)$. Finally, if $\Psi = \Phi$ or $\Psi = \psi \equiv \psi_L + \psi_R$, then

$$A_\alpha = \frac{1}{\sqrt{6}} (e_{4\alpha} + e_{\alpha 4}), \quad A_{\alpha+3} = \frac{i}{\sqrt{6}} (e_{4\alpha} - e_{\alpha 4}), \quad (17)$$

$$A_7 = \frac{1}{6} (e_{11} + e_{22} + e_{33} - 3e_{44}), \quad (18)$$

and A_a are defined by (16) as $a = 8, \dots, 15$. Here e_{ij} are matrices with the elements $(e_{ij})_{mn} = \delta_{im} \delta_{jn}$ and $\alpha = 1, 2, 3$.

It follows from (15) and (16) that the multiplet (8) must be $SU(3)$ -invariant. On the other hand, the skew-symmetric representation of $SU(4)$ has the $SU(3)$ content $\mathbf{6} = \mathbf{3} + \mathbf{3}^*$. Therefore in (13) and (14) we use the representation

$$\chi = \frac{1}{\sqrt{2}} [(u^{c1} e_{[23]} + u^{c2} e_{[31]} + u^{c3} e_{[12]})_L + (d^{c1} e_{[23]} + d^{c2} e_{[31]} + d^{c3} e_{[12]})_R], \quad (19)$$

where $e_{[ij]} = e_{ij} - e_{ji}$, in place of (8). Here we take advantage of the charge conjugation formula and $SU(4)$ -invariance of the Levi-Civita tensor ε^{ijkl} .

The hypercharge Y is equal to the doubled mean charge of the multiplet

$$Y = 2\langle Q \rangle, \quad (20)$$

and also

$$Y_\psi = 2Q - A_7. \quad (21)$$

It follows from (14) that

$$Q(\Phi_\alpha) = 1/3, \quad Q(\Phi_4) = 0. \quad (22)$$

Finally, the complex multiplet

$$\Phi^{ijk} = \frac{1}{\sqrt{2}} \varepsilon^{ijkl} \Phi_l. \quad (23)$$

For $\mu^2, \lambda > 0$, the vacuum expectation values of the components of Φ are

$$\langle \Phi_\alpha \rangle = 0, \quad \langle \Phi_0 \rangle \equiv v = \frac{\mu}{\sqrt{\lambda}}. \quad (24)$$

so that we have the breaking pattern

$$SU(4) \times U(1)_Y \rightarrow SU(3) \times U(1), \quad (25)$$

In order that to define the mass spectrum we define the fields

$$W_\mu'^\alpha = \frac{1}{\sqrt{2}} (A_\mu^\alpha - i A_\mu^{\alpha+3}) \quad (26)$$

and diagonalize the mass matrix by the orthogonal transformation

$$Z'_\mu = A_\mu^7 \cos \theta - B_\mu \sin \theta, \quad (27)$$

$$A_\mu = A_\mu^7 \sin \theta + B_\mu \cos \theta \quad (28)$$

with

$$\tan \theta = \frac{g'}{g}. \quad (29)$$

Then we get

$$M_{W'}^2 = \frac{g^2 v^2}{24}, \quad M_{Z'}^2 = \frac{v^2 (g^2 + g'^2)}{16}. \quad (30)$$

Substituting the fields (26)–(28) in the Lagrangian (9), we get

$$\mathcal{L} = \mathcal{L}_{QCD} + \mathcal{L}_{CC} + \mathcal{L}_{NC} + \dots, \quad (31)$$

where

$$\mathcal{L}_{QCD} = -\frac{1}{4} F_{\mu\nu}^k F^{k\mu\nu} + \bar{\psi}_q^{c\alpha} \left(i \gamma^\mu \tilde{D}_\mu + m_u \right) \psi_q^{c\alpha}. \quad (32)$$

Here the anti-quarks $u^{c\alpha}$ and $d^{c\alpha}$ are denoted by the symbols $\psi_q^{c\alpha}$,

$$\tilde{D}_\mu = \partial_\mu - ig \frac{\lambda_s}{2\sqrt{6}} A_\mu^s, \quad m_\psi = \frac{v}{\sqrt{2}} f_\psi. \quad (33)$$

Suppose m_ψ is the mass of ψ and g is connected with the effective QCD coupling α_s by the relation

$$g^2 = 24\pi\alpha_s. \quad (34)$$

Then $\mathcal{L}_{QCD}$ is exactly coincides with the Lagrangian describing the strong interactions of colored anti-quarks and gluons in the SM. The interaction of charged currents is described by the Lagrangian

$$\mathcal{L}_{CC} = -\frac{g}{2\sqrt{3}}J_\mu^\alpha W_\mu'^\alpha + h.c., \quad (35)$$

where

$$J_\mu^\alpha = \bar{d}_L^{\alpha\alpha}\gamma_\mu\nu_L + \bar{u}_R^{\alpha\alpha}\gamma_\mu e_R. \quad (36)$$

Obviously, the vector leptoquark $W_\mu'^\alpha$ has the electric charge $+1/3$. For the neutral currents in (31), we have

$$\mathcal{L}_{NC} = g\sin\theta J_{em}^\mu A_\mu + \frac{g}{\cos\theta}J_0^\mu Z_\mu. \quad (37)$$

where

$$J_{em}^\mu = Q^{(q)}\bar{\psi}_q^{c\alpha}\gamma^\mu\psi_q^{c\alpha} + Q^{(l)}\bar{\psi}_l\gamma^\mu\psi_l, \quad (38)$$

and

$$J_0^\mu = N_L^{(q)}\bar{\psi}_{qL}^{c\alpha}\gamma^\mu\psi_{qL}^{c\alpha} + N_R^{(q)}\bar{\psi}_{qR}^{c\alpha}\gamma^\mu\psi_{qR}^{c\alpha} + N_L^{(l)}\bar{\psi}_{lL}\gamma^\mu\psi_{lL} + N_R^{(l)}\bar{\psi}_{lR}\gamma^\mu\psi_{lR} \quad (39)$$

Here the anti-quarks and leptons are denoted by the symbols $\psi_q^{c\alpha}$ and ψ_l respectively, $Q^{(f)}$ is a charge of the fermion ψ_f and $N^{(f)}$ is a diagonal element of the matrix

$$N = \frac{A_7}{2} - Q\sin^2\theta. \quad (40)$$

Suppose that g is connected with the effective QED coupling α by the relation

$$g^2\sin^2\theta = 4\pi\alpha. \quad (41)$$

Then the first term in the right hand side of (37) is exactly coincides with the Lagrangian describing the electromagnetic interaction in the Glashow-Weinberg-Salam model. Comparing (34) and (41), we get

$$\sin^2\theta = \frac{\alpha}{6\alpha_s}. \quad (42)$$

Now we again imbed the $SU(4) \times U(1)_Y$ gauge group in the $SU(5)$ gauge group and consider the evolution of the coupling constants g and g' . The evolution of couplings in gauge theories with the groups $SU(n)$ is described by the renormalization group equation

$$\frac{dg_n}{d(\ln\mu)} = -b_n g_n^3. \quad (43)$$

Solution of this equation for g_n is

$$g_n^{-1}(\mu) = g_n^{-1}(\mu_0) - \frac{b_n}{8\pi^2} \ln \frac{\mu_0}{\mu}. \quad (44)$$

Since the $SU(4) \times U(1)_Y$ gauge theory has four multiplets of chiral fermions and one quartet of charged scalars, it follows that the one-loop beta function coefficients

$$b_4 = 44/3 - 4n/3 - 1/6, \quad (45)$$

$$b_1 = -20n/9 - 1/6, \quad (46)$$

where n is the number of generations. On the other hand, the embedding of $SU(4) \times U(1)_Y$ in the $SU(5)$ gauge group imposes the coupling constant relations

$$g^2 = 15g'^2 \quad (47)$$

as $\mu = \mu_0$. Using (29) and (41), we represent the relation (44) for $n = 1$ and 4 in the form

$$\alpha^{-1}(\mu)\sin^2\theta = \alpha_0^{-1} - \frac{1}{2\pi}b_4\ln\frac{\mu_0}{\mu} \quad (48)$$

$$\frac{1}{15}\alpha^{-1}(\mu)\cos^2\theta = \alpha_0^{-1} - \frac{1}{2\pi}b'_1\ln\frac{\mu_0}{\mu}, \quad (49)$$

where $b'_1 = b_1/15$. Given the experimental inputs [6]:

$$\alpha^{-1}(M_Z) = 127.909 \pm 0.019, \quad (50)$$

$$\alpha_s(M_Z) = 0.1217 \pm 0.017, \quad (51)$$

$$M_Z = 91.1874 \pm 0.0021 \text{ GeV} \quad (52)$$

we have

$$\sin^2\theta = 0.01071 \pm 0.00015, \quad (53)$$

$$\mu_0 = 5248 \pm 65 \text{ GeV}. \quad (54)$$

We assume that (54) defines the symmetry-breaking scale of the $SU(4) \times U(1)_Y$ gauge theory, i.e. that μ_0 is coincided with the vacuum expectation value v in (24). Then we get

$$M_{W'} = 3245 \pm 63 \text{ GeV}, \quad (55)$$

$$M_{Z'} = 3996 \pm 77 \text{ GeV}. \quad (56)$$

Further, we consider the vector current effective Lagrangian

$$\mathcal{L}_{eff} = -\frac{g}{12M_{W'}^2}J_\mu^\alpha \bar{J}^{\alpha\mu}, \quad (57)$$

in the limit of energies low compared to the W' mass. Suppose that

$$\frac{g^2}{12M_{W'}^2} = |\varepsilon|\frac{G_F}{\sqrt{2}}, \quad (58)$$

where G_F is the conventional Fermi coupling constant, known from the muon decay rate to have the value

$$G_F = 1.166367(5) \cdot 10^{-5} \text{ GeV}^{-2}. \quad (59)$$

Substituting (55) and (59) in (58), we find the factor

$$|\varepsilon| = (2.205 \pm 0.116) \times 10^{-3}. \quad (60)$$

It is interesting that this value is coincided with the measured CP-violating quantity $|\varepsilon_K|$. Note also that the CP-noninvariant interaction with coupling constants $G' \sim 10^{-3}G_F$ has been investigated in [7].

Now we consider the $SU(4) \times U(1)$ model with several generations. We rewrite the current (36) in the form

$$J_\mu^\alpha = \bar{\psi}_{qL}^{c\alpha} \gamma_\mu V'^{ql} \psi_{lL} + \bar{\psi}_{qR}^{c\alpha} \gamma_\mu V''^{ql} \psi_{lR}, \quad (61)$$

where V and V' are unitary mixing matrices. Further, suppose the generations of the $SU(4) \times U(1)$ model and CM are coincided. The current that couples to the W boson field in the CM is

$$J_\mu = \bar{\psi}_{qL}^\alpha \gamma^\mu V^{qq'} \psi_{q'L}^\alpha + \bar{\psi}_{lL} \gamma^\mu \psi_{lL}. \quad (62)$$

Comparing (61) and (62), we get either $V' = V^+$ and $V'' = 1$ or $V'' = V^*$ and $V' = 1$. We consider the case of two generations. In this case, V is a 2×2 unitary matrix. Such a matrix has 4 parameters: one is a rotation angle, and the other three are phases. We can remove two phases by performing the change of variables on the quark fields. Since the currents (61) and (62) contain both the particles and antiparticles, we cannot remove

the third phase. The same set of arguments can be made for the theory with three generations. The final form of the matrix V contains 3 angles and 3 phase. In particular, it follows from here that the Cabibbo-Kobayashi-Maskawa matrix [8] is obtained from V as a special case. Hence, the $SU(4) \times U(1)$ model can provide an alternative description of the observed CP violation.

* Electronic address: ek.loginov@mail.ru

- [1] Pati J. C., Salam A. Phys. Rev. **D10**, 275 (1974).
- [2] R. N. Mohapatra, J. C. Pati Phys. Rev. **D11**, 566 (1975); and G. Senjanovic, R. N. Mohapatra Phys. Rev. **D12**, 1502 (1975).
- [3] M. B. Voloshin, Sov. J. Nucl. Phys. **48**, 512 (1988); R. Foot, H. N. Long, and T. A. Tran, Phys. Rev. **D50**, 34 (1994); F. Pisano, V. Pleitez, Phys. Rev. **D51** 3865 (1995); D. E. Kaplan and M. Schmaltz, JHEP **0310**, 039 (2003).
- [4] L.-F. Li, Phys. Rev. **D9**, 1723 (1974).
- [5] H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
- [6] C. Amsler et al., Phys. Lett. **B667**, 1 (2008).
- [7] L. Wolfenstein, Phys. Rev. Lett. **13**, 562 (1964).
- [8] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).